

Reference U

RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 11 OF 25 HCPLUS COPYRIGHT 2008 ACS on STN
ACCESSION NUMBER: 2001:658540 HCPLUS
DOCUMENT NUMBER: 135:371618
TITLE: Isoquinoline syntheses via Δ^2 -oxazolines. Part
VIII. Cyclization of
2-acetamido-1,2-diphenylethan-1-ol derivatives into
isoquinoline systems
AUTHOR(S): Kopczynski, T.; Voeikel, A.
CORPORATE SOURCE: Institute of Chemical Technology and Engineering,
Poznan Technical University, Poznan, 60-965, Pol.
SOURCE: Polish Journal of Chemistry (2001), 75(9), 1317-1325
CODEN: PJCHDQ; ISSN: 0137-5083
PUBLISHER: Polish Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English
OTHER SOURCE(S): CASREACT 135:371618
AB The results of the conversion of 2-acetamido-1,2-diphenylethan-1-ol
derivs. into 1-methyl-4-phenylisoquinoline derivs. were described. The
mechanism proposed for these reaction assumes the existence of protonated
 Δ^2 -oxazolines, carbonium ions, and unsatd. amides as intermediates.
For example, the cyclization of erythro-N-(2-hydroxy-1,2-
diphenylethyl)acetamide or threo-N-(2-hydroxy-1,2-diphenylethyl)acetamide
gave 1-methyl-4-phenylisoquinoline in 66% yield.
IT 374594-09-9P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of isoquinolines via cyclocondensation of
N-(hydroxydiphenylethyl)acetamide derivs.)
RN 374594-09-9 HCPLUS
CN Isoquinoline, 6-methoxy-1-methyl-4-phenyl- (CA INDEX NAME)

